

Confronting pNGB quintessence with data

Koushik Dutta* and Lorenzo Sorbo†

Department of Physics, University of Massachusetts, Amherst, MA 01003

(Dated: February 5, 2008)

We analyze the observational constraints on the model where a pseudo-Nambu-Goldstone boson (pNGB) plays the role of dark energy. The constraints are derived by using the latest Gold set of 182 type Ia supernovae and the CMB shift parameter. We allow for both the initial value of the scalar field and the present value of the energy density in the pNGB to vary. We find that – compared to previous analyses – the allowed portion of parameter space has shrunk around the region where the pNGB does not evolve significantly.

I. INTRODUCTION

Since the type Ia supernova observations of [1, 2, 3], an intense activity has been devoted to the search of an explanation of the accelerated expansion of the Universe. If gravity is described by Einstein’s General Relativity and the effects of inhomogeneities can be neglected, then acceleration must be due to a dark energy component that represents roughly 70% of the matter content of the Universe. Current data tell that the equation of state parameter $w = p/\rho$ of dark energy has to obey $w \lesssim -0.7$ [4].

The simplest explanation of cosmic acceleration is a cosmological constant, the energy of vacuum, with magnitude

$$\Lambda \simeq (2 \times 10^{-3} \text{ eV})^4 \quad (1)$$

and equation of state parameter $w = -1$. This solution is attractive in many respects, both for its simplicity (a single parameter is needed to describe it) and for its excellent agreement with data. It is however hard to justify from a theoretical standpoint. Quantum fluctuations of matter, indeed, give contributions to the vacuum energy, and very precise cancellations are needed to keep this energy at small values. Unfortunately the Standard Model does not display any of these cancellations at least up to the scales that have been probed in collider experiments, about 60 orders of magnitude beyond the value of (1).

For this reason, soon after the release of [1, 2, 3], people have started to look for alternative scenarios, and a wide interest in *quintessence* models has emerged [5] (for a recent review, see [6]). The philosophy behind quintessence is the following. First, it is assumed that some mechanism is able to fix the energy of the ground state of the

Universe to zero¹. Then the existence of a new degree of freedom (quintessence) is postulated: quintessence is supposed not to have yet relaxed to its vacuum, so that its energy density is responsible for cosmic acceleration. Quintessence has $w \neq -1$ as a distinctive prediction, and is usually described by some scalar degree of freedom ϕ endowed with some potential $V(\phi)$. $V(\phi)$ has to be very flat, if we want w to be sufficiently negative.

It is possible to write down a virtually infinite number of quintessence potentials $V(\phi)$. However, only for few of them the flatness of the potential is not spoiled by radiative corrections and the exchange of quanta of ϕ do not give rise to an (unobserved) fifth force [7]. Those few potentials are more motivated from a theoretical point of view than the others. This is especially true for the pseudo-Nambu-Goldstone boson (pNGB) potential of [8] that has all the good qualities of radiative stability that anybody who believes in quantum mechanics might require.

In the present paper we analyze the parameter space of pNGB quintessence [8] in the light of the most recent observations, in particular those from supernovae. Our approach is orthogonal to the “model independent” approach recently taken on the subject by many investigators (see for instance [9]), and is admittedly based on a theoretical prejudice in favor of radiatively stable potentials. To our knowledge, the most recent complete analyses of this model date back to about five years ago [10, 11, 12]. Given the recent developments of the observational situation, we believe that it is important

¹ It is often stated that finding a mechanism that fixes to cosmological constant to zero should be easier than finding a mechanism that fixes it to some very small nonvanishing value. Let us note here that this is not what usually occurs in Quantum Field Theory: if it is possible to find a symmetry that fixes some quantity to zero, it is typically straightforward to break such a symmetry so that this quantity can be kept small in a controlled way.

*koushik@physics.umass.edu

†sorbo@physics.umass.edu

to perform an analysis of the model in which the latest data are taken into account.

In the next section we briefly describe the properties of the model of pNGB quintessence. Then in section III we present the observational constraints from supernovae and from the CMB shift parameter. In section IV we discuss our results before concluding in section V.

II. THE PNGB POTENTIAL

The use of pNGBs has been first proposed in order to realize a technically natural model of inflation in [13] and has been subsequently considered for dark energy in [8]. The model is characterized by a pseudoscalar field ϕ with a potential that can be well approximated by

$$V(\phi) = \mu^4 [\cos(\phi/f) + 1], \quad (2)$$

where we have neglected the contributions by higher harmonics (this is supposed to be a good approximation as long as f is sufficiently smaller than the Planck mass [14]). The potential is generated by the breaking of a shift symmetry $\phi \rightarrow \phi + \text{constant}$, and for this reason it is radiatively stable.

The cosmological evolution of this model is in general determined by four parameters: the quantities μ and f and the initial conditions ϕ_{in} and $\dot{\phi}_{\text{in}}$. Due to the high expansion rate of the Early Universe, we assume $\dot{\phi}_{\text{in}} = 0$. One more free parameter is eliminated if we insist that Ω_ϕ (the current ratio of the amount of dark energy over critical energy) is equal to 0.7. As a consequence, if we assume $\Omega_\phi = 0.7$, the model is described by only two parameters that can be taken to be f and ϕ_{in} . A detailed analysis of the dynamics of the pNGB zero mode can be found in [11]. Due to periodicity of the potential ϕ_{in} takes values between 0 and $2\pi f$. In addition, if we take into account the indication from String Theory [14] that f cannot be larger than $M_P \simeq 2.4 \times 10^{18}$ GeV, then the parameter space of the potential turns out to be compact. This implies that, at least in principle, all of it can be excluded by observation, and that pNGB quintessence in its simplest version can be ruled out. For this reason we find this model even more attractive (although there are ways to evade the constraint $f < M_P$ [15]), and we believe this is an additional motivation for studying it in detail. Also, for this reason we will restrict our study to the region $f < M_P$.

III. OBSERVATIONAL CONSTRAINTS

We consider the quintessence field ϕ with potential (2) in a flat Friedmann-Robertson-Walker Universe. The equations of motion are given by

$$H^2 = \frac{1}{3M_P^2} \left(\frac{1}{2} \dot{\phi}^2 + V(\phi) + \rho_m \right),$$

$$\ddot{\phi} + 3H\dot{\phi} + \frac{dV}{d\phi} = 0, \quad (3)$$

where $H = \dot{a}/a$ is the Hubble constant, ρ_m is the energy density of nonrelativistic matter and, since we will be dealing only with the dynamics of the late Universe, we neglect contributions by radiation. We solve these equations numerically to find the evolution of the scale factor as a function of time.

A. Type Ia supernovae

First, we investigate constraints from the observation of type Ia supernova from the data set [16], which is a compilation of old data [3] by HZS team, first year SuperNova Legacy Survey data [17] and recent observations of 21 new supernovae [4]. For our analysis we will consider only the 182 “high confidence” Gold SN data with $z > 0.0233$. Although most of the SNe have $z < 1$, there are 16 SNe with $z > 1$. This is the most up to date supernova data set available in the literature. This data set has been recently used in [18, 19] to study the observational constraints on different parametrizations of dark energy.

For a particular cosmological model with parameters $\mathbf{s}$ the predicted distance moduli are given by

$$\mu_0(z, \mathbf{s}) = m - M = 5 \log_{10} \left(\frac{d_L(\mathbf{s})}{\text{Mpc}} \right) + 25, \quad (4)$$

where m and M are the apparent and absolute magnitude of distant supernovae. d_L is the luminosity distance given by

$$d_L(z) = (1+z) \int_0^z \frac{dz'}{H(z')} \quad (5)$$

and depends only on the expansion history of the Universe from redshift z to today. Assuming that all the distance moduli are independent and normally distributed the likelihood function can be calculated from the chi-square statistics $\mathcal{L} \propto \exp(-\chi^2/2)$, where

$$\chi^2(\phi_{\text{in}}, \dot{\phi}_{\text{in}}, f, \mu, H_0) = \sum_{i=1}^{182} \frac{(\mu_{0i}^{\text{obs}} - \mu_{0i}^{\text{th}})^2}{\sigma_{0i}^2}. \quad (6)$$

Here μ_{0i}^{obs} and σ_{0i} are the measured value of the distance modulus and the corresponding uncertainty for the i -th supernova. μ_{0i}^{obs} and σ_{0i} , as well as the redshift z_i are found from the data set [16]. μ_0^{th} is calculated by using eq. (5), where $H(z)$ is obtained by numerically solving the background evolution equation (3). We marginalize the likelihood over the nuisance parameter H_0 [2].

As we have noted earlier, the model has four parameters: f, μ, ϕ_{in} and $\dot{\phi}_{in}$. We assume $\dot{\phi}_{in} = 0$ and allow the system to evolve until $\Omega_\phi = 0.7$ today. This leaves us with two parameters and we choose them to be f and ϕ_{in} . We plot the resulting confidence contour in figure 1. The upper left portion of the plot corresponds to the part of parameter space where Ω_ϕ does not reach the value 0.7. In this part of the parameter space, the scalar field rolls quickly to the minimum and oscillates around it, behaving like matter.




FIG. 1: The shaded areas at the bottom right of the figure denote the 1σ and 2σ confidence level regions for $\Omega_\phi = 0.7$. The upper left part of the plot corresponds to parameters for which Ω_ϕ never reaches 0.7. The white dashed line corresponds to the value of the parameters for which $w_0 = -0.965$.

The dark areas at the bottom right part of the plot give the 68.3% (1σ) and the 95.4% (2σ) confidence level regions. The 3σ contour runs between the 2σ contour and the boundary of the forbidden region (we do not show it in the plot for clarity). As we go closer to the boundary of the forbidden region, the value of χ^2 increases sharply. Even if the 95.4% confidence level area seems to cover almost all of the allowed region, this is actually not the case. Indeed, there is a part of the parameter space, below the boundary of the forbidden region, where Ω_ϕ goes across 0.7 several times, as an effect of the oscillations of ϕ . This means that points with the same values of ϕ_{in}, f

and Ω_ϕ can correspond to different histories, depending on the number of times Ω_ϕ has gone across the value 0.7. We have computed the value of χ^2 in the case where Ω_ϕ has crossed $\Omega_\phi = 0.7$ more than once and we have found that this part of the parameter space is excluded at more than the 3σ level.

Once $\Omega_\phi = 0.7$ is fixed, the constraints on the parameters ϕ_{in} and f can be converted into constraints on the plane (f, μ) . We plot the 1, 2, and 3σ contours in figure 2.




FIG. 2: 1, 2, and 3σ constraints on the plane (f, μ) for $\Omega_\phi = 0.7$. The lower part of the plot corresponds to values of the parameters for which Ω_ϕ cannot reach the value 0.7.

The thick white dashed line in the bottom right part of figure 1 corresponds to the part of parameter space that gives $w_0 = -0.965$ (where w_0 is the current value of the equation of state parameter). According to [20], $w < -0.965$ is the most optimistic constraint (at the 95.4% level) that we might obtain from future observations, should they converge to the regime where dark energy shows no evolution. Therefore the dashed line in figure 1 gives the most stringent 2σ constraint that we can expect to put on the parameters of pNGB quintessence.

We have also considered the case where the value of Ω_ϕ is allowed to vary. In this case we have fixed $f = M_P$ while keeping ϕ_{in} variable. In the left panel of figure 3 we show the 1, 2, and 3σ contours related to supernova observations on the (Ω_ϕ, ϕ_{in}) plane. The shaded upper right part of the plot is excluded since the corresponding value of Ω_ϕ cannot be reached. The contours are essentially vertical and centered around $\Omega_\phi \simeq 0.67$. However, at larger values of Ω_ϕ , somehow larger values of ϕ_{in} (cor-

responding to some evolution in the quintessence field) are allowed. The best fit is at $\Omega_\phi = 0.67$, $\phi_{\text{in}} = 0$ (so that the pNGB sits at the top of its potential and behaves as a cosmological constant) with $\chi^2 = 159.6$ for 180 degrees of freedom². For smaller values of f (figure not shown), the contours have the same shape, although they shrink along the ϕ_{in} direction.




FIG. 3: Constraints on the pNGB parameter space for $f = M_P$. Left panel: 1, 2 and 3σ confidence level contours from supernovae only (dashed lines) and from the CMB shift parameter (7) only (solid lines). Right panel: 1, 2 and 3σ contours from the joint analysis.

B. CMB shift parameter

In addition to the SN data, we use Cosmic Microwave Background (CMB) data to constrain the model. In particular we derive constraints from the CMB shift parameter $\mathcal{R}$, that measures the shift in the angular size of the acoustic peaks of CMB when parameters of the theory are varied. $\mathcal{R}$ is independent on the present value of the Hubble constant, and is given by

$$\mathcal{R} = \sqrt{\Omega_m} H_0 \int_0^{z_{\text{cmb}}} \frac{dz}{H(z)}, \quad (7)$$

where z_{cmb} is the redshift of recombination. By using WMAP 3rd year data the value of the shift parameter has been extracted as $\mathcal{R} = 1.70 \pm 0.03$ for $z_{\text{cmb}} = 1089$ [21]. In

the left panel of figure 3 we plot the resulting confidence contour arising from this constraint. Note that the CMB contours favor a value of Ω_ϕ that is slightly larger than that favored by the SN data.

Once the SN and the CMB constraints are combined, we obtain the plot shown on the right panel of figure 3. Since we are assuming a flat Universe, imposing the shift parameter constraint does not reduce significantly the area of the allowed region, and indeed the SN and CMB constraints are not orthogonal. However, the CMB constraint helps eliminate the part of parameter space at large Ω_ϕ and large ϕ_{in} that is available at the 2σ and 3σ level if only SN constraints are taken into account. Moreover, once the CMB constraint is added, the best fit point is not any more at $\phi_{\text{in}} = 0$, but at the point $\phi_{\text{in}} = 1.25 M_P$, $\Omega_\phi = 0.71$ where $\chi^2 = 161.6$. This should be compared to $\chi^2 = 162.9$ found at $\Omega_\phi = 0.71$ when the constraint $\phi_{\text{in}} = 0$ is imposed. Since $\phi_{\text{in}} \neq 0$ implies that ϕ is rolling, the combination of CMB and SN data seems to hint at some evolution in dark energy. However, this hint should be taken with a grain of salt, since it emerges when we join two data sets that are not exactly compatible, as shown by the increase of ~ 2 units in χ^2 when we add the single CMB point to the SN data set (as stated above, the best fit point for SN data only has $\chi^2 = 159.6$).

IV. DISCUSSION

The data presented above indicate that, if we use supernova constraints only, the parameters that yield the best fit to data are those where the field ϕ sits at the top of the cosine potential, thus mimicking a cosmological constant. If we use also the constraint from the CMB shift parameter, a slowly rolling pNGB is slightly preferred to a constant one.

In order to see how the new data of [4] improve the constraints on the model, we can compare our results with those of previous analyses. In [10], Waga and Frieman have studied the constraints from the 1998 supernova data of Riess et al. [1], together with the statistics of gravitationally lensed quasars. Comparison of our results with those of [10] is complicated by different assumptions on the parametrization of the model. Indeed, in [10] the value of ϕ_{in} is fixed to $1.5 f$, so that Ω_ϕ is not a free parameter, but is function of f and μ . Nevertheless, some comparison is possible: in the parameter space of [10] there is still room at 2σ for a small region where the scale factor of the Universe is currently decelerating. In our analysis (see figure 4) this is not possible any more

² The best-fit χ^2 for the older dataset [3] was of $\chi^2 = 178.1$ for 155 degrees of freedom. The lower value of χ^2 for the current data set can be largely attributed to more conservative assumptions on the dispersions σ_{0i} .

at the 2σ level, even if it is still allowed at 3σ .

In [11], a wider portion of the parameter space is analyzed (and a different data set [2] is used), that shows that a part of parameter space where ϕ has performed half oscillation is allowed at the 2σ level. As we have stressed in the section III, the current data do not allow for this possibility any more.

In [12], a detailed study of the parameter space of the model has been performed by taking into account the constraints from CMB observations of BOOMERanG [22] and MAXIMA [23]. Constraints on this quintessence model were derived from its effects on integrated Sachs-Wolfe effect as well as from its effects on the location of the first peak. In figure 2 we show the $1, 2$, and 3σ constraints in the (f, μ) plane obtained by our analysis. Comparison with figure 5 of [12], shows that the more recent data improve by a factor of three or so the constraints on μ .

Let us also note that a quintessence field that is climbing up the potential could mimic $w < -1$ [24, 25] and possibly offer a better fit to data (see however [26]). In our case this is possible only if Ω_ϕ has already gone through a maximum. As we have mentioned above, this case appears to be not realized at the 3σ confidence level for a cosine potential. Indeed, a more asymmetric potential (such as that pictured in figure 1 of [25]) is needed to impart a sufficiently large velocity to the field and improve the fit to data.




FIG. 4: $1, 2$, and 3σ contours in the plane (Ω_{DE}, w_0) for the pNGB model with $f = M_P$ (dashed, thicker lines) and for dark energy with constant equation of state (solid lines).

Finally, it is instructive to discuss the current value of the equation of state parameter w_0 as obtained in the pNGB model and to compare it with the value of w_0 obtained by assuming that it is constant throughout the

evolution of the Universe. In figure 4 we show the plots of the $1, 2$ and 3σ confidence level contours in the plane (Ω_ϕ, w_0) both for the case of pNGB quintessence with $f = M_P$ and for a model with constant w_0 (only supernova data are used to compute the contours in figure 4). In the case of constant w_0 the contours are quite tilted, allowing for a value of w_0 significantly different from -1 only if Ω_{DE} gets very close to unity. Indeed, if dark energy has a constant equation of state parameter different from -1 , then a larger amount of dark energy is needed to get the same averaged value of w . In the case of a pNGB, however, the value of w_0 can be significantly different from -1 even if w was close to -1 in the past. As a consequence, one can obtain the required averaged value of w even without requiring that Ω_ϕ is extremely close to unity.

V. CONCLUSIONS

We have analyzed the portion of parameter space available for the model of pNGB quintessence. Our work extends the previous studies on the subject by allowing both for variations in the initial value of the zero mode of the pNGB and for variations in the current value of Ω_ϕ . Using the most up-to-date supernova data, we have shown that the parameter space of the pNGB potential is significantly constrained around the region where quintessence is sitting on the top of the cosine potential or slowly evolving along it. At the 95.4% level, previous analyses on the subject [10, 11] were still allowing the current value of w to be larger than $-1/3$ or even the possibility that quintessence had already performed a half oscillation about its minimum. Current data do not allow this any more.

We have also observed that, when CMB and SN constraints are joined, an evolving pNGB provides a slightly better fit to data than a pNGB stuck at the top of its potential.

Let us finally discuss future perspectives. Already now, data tell that f cannot be smaller than about a third of Planck mass (unless we fine tune ϕ_{in} to be very close to the top of the potential). As shown in figure 1, future data might constrain $f \gtrsim M_P/2$, leading to some tension with the requirement $f \lesssim M_P$ from String Theory [14]. One might wonder if this will be enough to consider the model “finely tuned”, and to start to consider alternative options [15] as more natural. But one can also take a more optimistic approach: maybe future data will show that cosmic acceleration is sourced by an *evolving*, radiatively stable pseudo-Nambu-Goldstone boson.

Acknowledgments

We thank Subinoy Das and John Donoghue for useful discussions.

-
- [1] A. G. Riess *et al.*, *Astron. J.* **116** (1998) 1009. [astro-ph/9805201].
 - [2] S. Perlmutter *et al.*, *Astrophys. J.* **517** (1999) 565 [astro-ph/9812133]; J. L. Tonry *et al.*, *Astrophys. J.* **594** (2003) 1 [arXiv:astro-ph/0305008]; R. A. Knop *et al.* [Supernova Cosmology Project Collaboration], *Astrophys. J.* **598**, 102 (2003) [arXiv:astro-ph/0309368].
 - [3] A. G. Riess *et al.* [Supernova Search Team Collaboration], *Astrophys. J.* **607**, 665 (2004) [arXiv:astro-ph/0402512].
 - [4] A. G. Riess *et al.*, arXiv:astro-ph/0611572.
 - [5] C. Wetterich, *Nucl. Phys. B* **302**, 668 (1988). B. Ratra and P. J. E. Peebles, *Phys. Rev. D* **37**, 3406 (1988). R. R. Caldwell, R. Dave and P. J. Steinhardt, *Phys. Rev. Lett.* **80**, 1582 (1998) [arXiv:astro-ph/9708069].
 - [6] E. J. Copeland, M. Sami and S. Tsujikawa, arXiv:hep-th/0603057.
 - [7] S. M. Carroll, *Phys. Rev. Lett.* **81** (1998) 3067 [arXiv:astro-ph/9806099]; C. F. Kolda and D. H. Lyth, *Phys. Lett. B* **458** (1999) 197 [arXiv:hep-ph/9811375]; T. Chiba, *Phys. Rev. D* **60** (1999) 083508 [arXiv:gr-qc/9903094]; P. Brax and J. Martin, *Phys. Lett. B* **468**, 40 (1999) [arXiv:astro-ph/9905040].
 - [8] J. A. Frieman, C. T. Hill, A. Stebbins and I. Waga, *Phys. Rev. Lett.* **75** (1995) 2077 [arXiv:astro-ph/9505060].
 - [9] J. Weller and A. Albrecht, *Phys. Rev. D* **65**, 103512 (2002) [arXiv:astro-ph/0106079]; V. Sahni, T. D. Saini, A. A. Starobinsky and U. Alam, *JETP Lett.* **77**, 201 (2003) [Pisma Zh. Eksp. Teor. Fiz. **77**, 249 (2003)] [arXiv:astro-ph/0201498]; U. Alam, V. Sahni, T. D. Saini and A. A. Starobinsky, *Mon. Not. Roy. Astron. Soc.* **344**, 1057 (2003) [arXiv:astro-ph/0303009]; U. Alam, V. Sahni, T. D. Saini and A. A. Starobinsky, *Mon. Not. Roy. Astron. Soc.* **344**, 1057 (2003) [arXiv:astro-ph/0303009]; Z. K. Guo, N. Ohta and Y. Z. Zhang, *Phys. Rev. D* **72**, 023504 (2005) [arXiv:astro-ph/0505253]. R. R. Caldwell and E. V. Linder, *Phys. Rev. Lett.* **95**, 141301 (2005) [arXiv:astro-ph/0505494]; Z. K. Guo, N. Ohta and Y. Z. Zhang, arXiv:astro-ph/0603109; V. Sahni and A. Starobinsky, arXiv:astro-ph/0610026; D. Huterer and H. V. Peiris, arXiv:astro-ph/0610427; C. Li, D. E. Holz and A. Cooray, arXiv:astro-ph/0611093.
 - [10] I. Waga and J. A. Frieman, *Phys. Rev. D* **62**, 043521 (2000) [arXiv:astro-ph/0001354].
 - [11] S. C. C. Ng and D. L. Wiltshire, *Phys. Rev. D* **63**, 023503 (2001) [arXiv:astro-ph/0004138].
 - [12] M. Kawasaki, T. Moroi and T. Takahashi, *Phys. Rev. D* **64**, 083009 (2001)
 - [13] K. Freese, J. A. Frieman and A. V. Olinto, *Phys. Rev. Lett.* **65** (1990) 3233; F. C. Adams, J. R. Bond, K. Freese, J. A. Frieman and A. V. Olinto, *Phys. Rev. D* **47**, 426 (1993) [arXiv:hep-ph/9207245].
 - [14] M. Dine, arXiv:hep-th/0107259; T. Banks, M. Dine, P. J. Fox and E. Gorbatov, *JCAP* **0306**, 001 (2003) [arXiv:hep-th/0303252].
 - [15] J. E. Kim and H. P. Nilles, *Phys. Lett. B* **553**, 1 (2003) [arXiv:hep-ph/0210402]; N. Arkani-Hamed, H. C. Cheng, P. Creminelli and L. Randall, *Phys. Rev. Lett.* **90** (2003) 221302; [arXiv:hep-th/0301218]; *JCAP* **0307** (2003) 003, [arXiv:hep-th/0302034]; L. Pilo, D. A. Rayner and A. Riotto, *Phys. Rev. D* **68** (2003) 043503. [arXiv:hep-ph/0302087]; J. E. Kim, H. P. Nilles and M. Peloso, *JCAP* **0501**, 005 (2005) [arXiv:hep-ph/0409138]; S. Dimopoulos, S. Kachru, J. McGreevy and J. G. Wacker, arXiv:hep-th/0507205; N. Kaloper and L. Sorbo, *JCAP* **0604**, 007 (2006) [arXiv:astro-ph/0511543].
 - [16] http://braeburn.pha.jhu.edu/~ariess/R06/sn_sample
 - [17] P. Astier *et al.*, *Astron. Astrophys.* **447**, 31 (2006) [arXiv:astro-ph/0510447].
 - [18] V. Barger, Y. Gao and D. Marfatia, arXiv:astro-ph/0611775.
 - [19] Y. G. Gong and A. z. Wang, arXiv:astro-ph/0612196.
 - [20] A. Albrecht *et al.*, arXiv:astro-ph/0609591.
 - [21] Y. Wang and P. Mukherjee, *Astrophys. J.* **650**, 1 (2006) [arXiv:astro-ph/0604051].
 - [22] P. de Bernardis *et al.* [Boomerang Collaboration], *Nature* **404**, 955 (2000) [arXiv:astro-ph/0004404].
 - [23] S. Hanany *et al.*, *Astrophys. J.* **545**, L5 (2000) [arXiv:astro-ph/0005123].
 - [24] M. Sahlen, A. R. Liddle and D. Parkinson, *Phys. Rev. D* **72**, 083511 (2005) [arXiv:astro-ph/0506696].
 - [25] C. Csaki, N. Kaloper and J. Terning, *JCAP* **0606**, 022 (2006) [arXiv:astro-ph/0507148];
 - [26] M. Sahlen, A. R. Liddle and D. Parkinson, arXiv:astro-ph/0610812.